

BCQ

Natural Anti-Inflammatory Agents

Monograph Prepared by Herb Joiner-Bey, N.D.

Boswellia serrata Extract (65% boswellic acids)	200mg
Bromelain (2400gdu-3600mcu/gm)	100mg
Curcumin Extract (95%)	200mg
Quercetin	100mg

Recommended dosage: 2 or 3 capsules 3-4 times per day.

INDICATIONS:

- Chronic inflammation
- Rheumatoid arthritis
- Sports injury
- Acute traumatic inflammation

Boswellia serrata

- Major constituents: boswellic acids (pentacyclic triterpenic acids) and essential oils
- Used traditionally in Indian Ayurvedic medicine to alleviate symptoms of chronic rheumatic inflammation.
- Inhibits the production of pro-inflammatory leukotrienes by the enzyme 5-lipoxygenase.
- Reduces the degradation of glycosaminoglycans, major components of connective tissue, in experimental models with lab animals. (Wildfeuer, Ammon, Satavhi)
- Three 400 mg tablets 2 or 3 times daily of Boswellia extract was found to be effective in the treatment 260 individuals with rheumatoid arthritis. (Etzel)

Bromelain

- Mixture of digestive enzymes from the stem of the pineapple plant (*Ananas comosus*).
- Reduces pain and inflammation associated with arthritis, trauma, or sports injury. (Klein)
- Prevents edema after traumatic incident or surgery.
- Prevents and reduces edema by inhibiting formation of fibrin within damaged tissue, allowing lymph fluids to pass with less obstruction.
- May interfere with the arachidonic acid cascade, thereby impeding the formation of pro-inflammatory eicosanoids. (Vellini)
- Blocks synthesis of kinin compounds that increase edema and pain. (Ito)

Curcumin

- Yellow pigment of turmeric rhizome (*Curcuma longa*).
- Acts as a powerful antioxidant against active oxygen moieties.
- Displays anti-inflammatory effects equivalent to some NSAIDs in acute models of inflammation and half as potent in chronic inflammation. (Satoskar)
- Inhibits lipoxygenase and cyclooxygenase and the formation of inflammatory leukotrienes and prostaglandins. (Huang)
- Promotes fibrinolysis.
- Inhibits the production of pro-inflammatory peroxynitrite radicals and nitrite. (Chan)
- Stabilizes lysosomal membranes. (Wichtl, p174)
- Inhibits pro-inflammatory gene expression. (Jobin)

Natural Anti-Inflammatory Agents (page 2)

Quercetin

- Backbone (aglycone) of a number of other flavonoid compounds.
- One of the most active flavonoid compounds studied.
- Inhibits manufacture and release of histamine from mast cells and basophils.
- Exhibits significant antioxidant activity, neutralizing free radical moieties, which can exacerbate inflammation.
- Demonstrates an inhibitory effect on acute inflammation in lab animal models. (Taguchi, Romero)

REFERENCES

1. Wildfeuer A, et al, Effects of boswellic acids extracted from an herbal medicine on the biosynthesis of leukotrienes and the course of experimental autoimmune encephalomyelitis. *Arzneimittelforschung* 1998 June;48(6):668-74.
2. Ammon HP, et al, Mechanism of anti-inflammatory actions of curcumin and boswellic acids. *J Ethnopharmacol* 1993 Mar;38(2-3):113-9.
3. Safayhi H, et al, Boswellic acids: novel. Specific, non-redox inhibitors of 5-lipoxygenase. *J Pharmacol Exp Ther* 1992 Jun;261(3):1143-6.
4. Klein G and Kullich W, Reducing pain by oral enzyme therapy in rheumatic diseases. *Wien Med Wochenschr* 1999;149(21-22):577-80.
5. Vellini M, et al, Possible involvement of eicosanoids in the pharmacological action of bromelain. *Arzneimittelforschung* 1986;36(1):110-2.
6. Ito C, et al, Anti-inflammatory actions of proteases, bromelain, trypsin and their mixed preparation. *Nippon Yakurigaku Zasshi* 1979 Apr 20;75(3):227-37.
7. Jobin C, et al, Curcumin blocks cytokine-mediated NF-kappa B activation and proinflammatory gene expression by inhibiting inhibitory factor I-kappa B kinase activity. *J Immunol* 1999 Sep 15;163(6):3474-83.
8. Satoskar RR, et al, Evaluation of anti-inflammatory property of curcumin (diferuloyl methane) in patients with postoperative inflammation. *Int J Clin Pharmacol Ther Toxicol* 1986 Dec;24(12):651-4.
9. Huang MT, et al, Inhibitory effects of curcumin on in vitro lipoxygenase and cyclooxygenase activities in mouse epidermis. *Cancer Res* 1991 Feb 1;51(3):813-9.
10. Chan MM, et al, Effects of three dietary phytochemicals from tea, rosemary and turmeric on inflammation-induced nitrite production. *Cancer Lett* 1995 Sep 4;96(1):23-9.
11. Cnandra D and Gupta S, Anti-inflammatory and anti-arthritis activity of a volatile oil of *Curcuma longa* (Haldi). *Indian J Med Res* 1972;60:138-142.
12. Arora R, et al, Anti-inflammatory studies on *Curcuma longa* (turmeric). *Indian J Med Res* 1971;59:1289-95.
13. Taguchi K, et al, Pharmacological studies of *Houttuynia herba*: the anti-inflammatory effect of quercitrin. *Yakugaku Zasshi* 1993 Apr;113(4):327-33.
14. Romero J, et al, Pharmacologic modulation of acute ocular inflammation with quercetin. *Ophthalmic Res* 1989;21(2):112-7.
15. Werbach and Murray M, *Botanical Influences on Illness: A Sourcebook of Clinical Research*. Tarzana, CA: Third Line Press, 1994.
16. Wichtl M, ed., *Herbal Drugs and Phytopharmaceuticals: A Handbook for Practice on a Scientific Basis*. Boca Raton: CRC Press, 1994.
17. Etzel, R., Special Extract of *Boswellia serrata* (H 15) in the Treatment of Rheumatoid Arthritis, *Phytomedicine*, 1996;3(1):91-94.